



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Ventricular rupture in Takotsubo cardiomyopathy**

Jaguszewski, M ; Fijalkowski, M ; Nowak, R ; Czapiewski, P ; Ghadri, J-R ; Templin, C ; Rynkiewicz, A

DOI: <https://doi.org/10.1093/eurheartj/ehs054>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-153781>

Journal Article

Published Version

Originally published at:

Jaguszewski, M; Fijalkowski, M; Nowak, R; Czapiewski, P; Ghadri, J-R; Templin, C; Rynkiewicz, A (2012). Ventricular rupture in Takotsubo cardiomyopathy. *European Heart Journal*, 33(8):1027.

DOI: <https://doi.org/10.1093/eurheartj/ehs054>

## CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehs054

Online publish-ahead-of-print 12 March 2012

## Ventricular rupture in Takotsubo cardiomyopathy

Milosz Jaguszewski<sup>1,2,\*†</sup>, Marcin Fijalkowski<sup>1</sup>, Radosław Nowak<sup>1</sup>, Piotr Czapiewski<sup>3</sup>, Jelena-Rima Ghadri<sup>2</sup>, Christian Templin<sup>2</sup>, and Andrzej Rynkiewicz<sup>1</sup>

<sup>1</sup>First Department of Cardiology, Medical University of Gdansk, Gdansk 80-952, Poland; <sup>2</sup>Department of Cardiology, Cardiovascular Center, University Hospital Zurich, Zurich, Switzerland; and <sup>3</sup>Department of Pathology, Medical University of Gdansk, Gdansk, Poland

\* Corresponding author. Tel: +41 44 255 11 11, Fax: +41 44 255 44 01, Email: [jamilosz@gmail.com](mailto:jamilosz@gmail.com), [milosz.jaguszewski@usz.ch](mailto:milosz.jaguszewski@usz.ch)

† Present address: Department of Cardiology, University Hospital Zurich, 8091 Zurich, Switzerland.

An 82-year-old female after a stress event, with no past medical history of cardiovascular diseases, was referred for emergency coronary angiography. She was suffering from chest pain, with a blood pressure of 124/67 mmHg and a heart rate of 76 b.p.m. Prominent ST-segment elevation in V1–V5 and increased troponin I level (14.82 ng/mL) suggested anteroapical acute myocardial infarction. Urgent angiography documented no coronary artery disease (Panel A). Bedside echocardiography revealed abnormal left ventricular (LV) contraction with an 'apical ballooning' pattern; LV ejection fraction was 55% (Panel B). The bull's-eye parametric image of the peak longitudinal systolic strain indicated dyskinesia of the apical segments, not pathognomonic for acute myocardial infarction (Panel B).

With subsequent diagnosis of Takotsubo cardiomyopathy (TTC), the patient remained asymptomatic. On the second day of hospitalization, a pseudoaneurysm in the inguinal area was documented. After unsuccessful treatment with freehand ultrasound-guided compression, the patient was referred for vascular surgery simultaneously receiving 6% hydroxyethyl starch infusion on the fifth day of hospitalization. During transportation to the operating room, electrocardiogram monitoring documented bradysystolia. The patient was promptly reanimated and urgent echocardiography revealed pericardial effusion with signs of tamponade. Despite extended and extensive resuscitation efforts, the patient died.

The autopsy identified a wide penetrating apical rupture as well as 1500 mL of thrombi and liquid blood in the pericardium (Panel C). A tissue obtained from the focus of the apical fissure revealed fields of hypertrophied and disarrayed cardiomyocytes, surrounded by predominantly mononuclear inflammatory infiltrate and loose connective tissue as well the foci of haemorrhage (Panel D). Myocarditis was excluded by the pathologist.

The presented case should elucidate that TTC is potentially life threatening in the acute phase, despite a large body of literature reporting that it is a benign disease in most cases.

